

Abnormal Structure of Fermion Mixings in a Seesaw Quark Mass Matrix Model

Yoshio Koide*

Department of Physics, University of Shizuoka
395 Yada, Shizuoka 422, Japan

Abstract

It is pointed out that in a seesaw quark mass matrix model which yields a singular enhancement of the top-quark mass, the right-handed fermion-mixing matrix U_R^u for the up-quark sector has a peculiar structure in contrast to the left-handed one U_L^u . As an example of the explicit structures of U_L^u and U_R^u , a case in which the heavy fermion mass matrix M_F is given by a form [(unit matrix)+(rank-one matrix)] is investigated. As a consequence, one finds observable signatures at projected high energy accelerators like the production of a fourth heavy quark family.

I. INTRODUCTION

Why is the top-quark mass m_t so singularly enhanced compared with the bottom-quark mass m_b (while keeping $m_u \sim m_d$)? Why does only the top-quark have a mass of the order of the electroweak scale m_W ? Recently, it has been pointed out [1,2] that a seesaw quark mass matrix model [3] can give a natural answer to these questions.

In the conventional seesaw mass matrix model [3], we assume vector-like heavy fermions F_i in addition to the conventional three-family quarks and leptons f_i ($f = u, d, \nu, e$; $i = 1, 2, 3$). These fermions f and F belong to $f_L = (2, 1)$, $f_R = (1, 2)$, $F_L = (1, 1)$ and $F_R = (1, 1)$ of $SU(2)_L \times SU(2)_R$. The fermions F acquire masses M_F at a large energy scale $\mu \sim \lambda m_0$. The symmetries $SU(2)_L$ and $SU(2)_R$ are broken at the energy scales $\mu \sim m_0$ and $\mu \sim \kappa m_0$, respectively. Then, the 6×6 mass matrix M for the fermions (f, F) is given by

$$M = \begin{pmatrix} 0 & m_L \\ m_R & M_F \end{pmatrix} = m_0 \begin{pmatrix} 0 & Z_L \\ \kappa Z_R & \lambda Y \end{pmatrix}, \quad (1.1)$$

where matrices Z_L , Z_R and Y are dimensionless matrices with the order of one. For $|\lambda| \gg |\kappa| \gg 1$ and $\det M_F \neq 0$, the 3×3 mass matrix for fermions f , M_f , is approximately given by the well-known ‘‘seesaw’’ expression [4]

$$M_f \simeq -m_L M_F^{-1} m_R, \quad (1.2)$$

so that the fermion masses m_i^f ($i = 1, 2, 3$) are suppressed by a factor κ/λ to the electroweak scale m_0 . This was one of the motivations for considering a seesaw mechanism for quarks [3] before the discovery of the top quark [5]. However, the observation of the large top-quark mass has demanded that top-quark mass should be of the order of m_0 without the factor κ/λ .

Recently, it has been found [1,2] that the seesaw mass matrix with $\det M_F = 0$ can yield fermion masses m_i^f and $m_i^F \equiv m_{i+3}^f$ ($i = 1, 2, 3$) with the following order:

$$\begin{aligned} m_1, m_2 &\sim (\kappa/\lambda)m_0, \\ m_3 &\sim m_0 \sim O(m_L), \\ m_4 &\sim \kappa m_0 \sim O(m_R), \\ m_5, m_6 &\sim \lambda m_0 \sim O(M_F). \end{aligned} \quad (1.3)$$

Note that the third fermion mass does not have the factor κ/λ . Therefore, if the heavy fermion mass matrix M_F takes $\det M_F = 0$ in the up-quark sector, we can

understand why only the top-quark has a mass of the order of m_L without the suppression factor κ/λ . This was first explicitly derived by Fusaoka and the author [1] on the basis of a special seesaw mass matrix model, “democratic seesaw mass matrix model”, where M_F is given by the form [(unit matrix)+(a rank-one matrix)], and then generalized by Morozumi *et al.* [2].

In the present paper, we will point out that in such a model the right-handed fermion-mixing matrix U_R has a peculiar structure in contrast to the left-handed one U_L , i.e., as if the third and fourth rows of U_R are exchanged each other in contrast to U_L . In Sec. II, we will discuss general properties of the fermion mass spectrum and the mixing matrices U_L and U_R in the would-be seesaw mass matrix (1.1) with $\det M_F = 0$. In Sec. III, in order to see the more explicit relations between the Cabibbo-Kobayashi-Maskawa (CKM) [6] matrices V_L and V_R , we investigate a case with constraints $Z_R^T = Z_L$ and $Y^T = Y$ which are not so restrictive and which most models can satisfy. In Sec. IV, as an explicit example of U_L and U_R , we evaluate the mixing matrices for the case of the “democratic seesaw mass matrix model” which can yield realistic quark masses and CKM matrix V_L . The final section V will be devoted to the summary.

II. GENERAL STRUCTURES OF THE FERMION MIXING MATRICES

The mixing matrices U_L and U_R are obtained by diagonalizing the following Hermitian matrices H_L and H_R , respectively:

$$\begin{aligned} H_L &\equiv MM^\dagger = \begin{pmatrix} m_L m_L^\dagger & m_L M_F^\dagger \\ M_F m_L^\dagger & M_F M_F^\dagger + m_R m_R^\dagger \end{pmatrix} \\ &= m_0^2 \begin{pmatrix} Z_L Z_L^\dagger & \lambda Z_L Y^\dagger \\ \lambda Y Z_L^\dagger & \lambda^2 Y Y^\dagger + \kappa^2 Z_R Z_R^\dagger \end{pmatrix}, \end{aligned} \quad (2.1)$$

$$\begin{aligned} H_R &\equiv M^\dagger M = \begin{pmatrix} m_R^\dagger m_R & m_R^\dagger M_F \\ M_F^\dagger m_R & M_F^\dagger M_F + m_L^\dagger m_L \end{pmatrix} \\ &= m_0^2 \begin{pmatrix} \kappa^2 Z_R^\dagger Z_R & \kappa \lambda Z_R^\dagger Y \\ \kappa \lambda Y^\dagger Z_R & \lambda^2 Y^\dagger Y + Z_L^\dagger Z_L \end{pmatrix}. \end{aligned} \quad (2.2)$$

For $\det M_F \neq 0$, we can obtain the well-known seesaw expression (1.2) since the

3×3 Hermitian matrices H_L^f and H_R^f for fermions f_i are approximately given by

$$H_L^f \simeq m_L m_L^\dagger - m_L M_F^\dagger (M_F M_F^\dagger + m_R m_R^\dagger)^{-1} M_F m_L^\dagger \simeq -m_L M_F^{-1} m_R (m_L M_F^{-1} m_R)^\dagger , \quad (2.3)$$

$$H_R^f \simeq m_R^\dagger m_R - m_R^\dagger M_F (M_F^\dagger M_R + m_L^\dagger m_L)^{-1} M_F^\dagger m_R \simeq -(m_L M_F^{-1} m_R)^\dagger m_L M_F^{-1} m_R . \quad (2.4)$$

The 6×6 mixing matrices U_L and U_R are approximately given by

$$U_L \simeq \begin{pmatrix} A_L & 0 \\ 0 & B_L \end{pmatrix} \begin{pmatrix} \mathbf{1} & -m_L M_F^{-1} \\ M_F^{\dagger-1} m_L^\dagger & \mathbf{1} \end{pmatrix} = \begin{pmatrix} A_L & -A_L m_L M_F^{-1} \\ B_L M_F^{\dagger-1} m_L^\dagger & B_L \end{pmatrix} , \quad (2.5)$$

$$U_R \simeq \begin{pmatrix} A_R & 0 \\ 0 & B_R \end{pmatrix} \begin{pmatrix} \mathbf{1} & -m_R^\dagger M_F^{\dagger-1} \\ M_F^{-1} m_R & \mathbf{1} \end{pmatrix} = \begin{pmatrix} A_R & -A_R m_R^\dagger M_F^{\dagger-1} \\ B_R M_F^{-1} m_R & B_R \end{pmatrix} , \quad (2.6)$$

where the 3×3 unitary matrices A and B are defined by

$$-A_L m_L M_F^{-1} m_R A_R^\dagger = D_f , \quad B_L M_F B_R^\dagger = D_F , \quad (2.7)$$

$D_f = \text{diag}(m_1^f, m_2^f, m_3^f)$ and $D_F = \text{diag}(m_4^f, m_5^f, m_6^f) \equiv \text{diag}(m_1^F, m_2^F, m_3^F)$. The mixing matrix U_R has a structure similar to U_L except for the point that the off-diagonal elements $(U_L)_{ik}$ and $(U_L)_{ki}$ ($i = 1, 2, 3$; $k = 4, 5, 6$) have a suppression factor $1/\lambda$, while $(U_R)_{ik}$ and $(U_R)_{ki}$ have a suppression factor κ/λ .

On the other hand, for the case of $\det M_F = 0$, the seesaw expression (1.2) is not valid any longer. For the case of $\det M_F = 0$, without losing generality, we can choose a heavy fermion basis where the mass matrix M_F is given by a diagonal form

$$M_F = \lambda m_0 \begin{pmatrix} 0 & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix} , \quad (2.8)$$

where $*$ denote elements with the order of one. Then the Hermitian matrices (2.1)

and (2.2) take the following textures:

$$H_L = m_0^2 \begin{pmatrix} * & * & * & 0 & \sim \lambda & \sim \lambda \\ * & * & * & 0 & \sim \lambda & \sim \lambda \\ * & * & * & 0 & \sim \lambda & \sim \lambda \\ 0 & 0 & 0 & \sim \kappa^2 & \sim \kappa^2 & \sim \kappa^2 \\ \sim \lambda & \sim \lambda & \sim \lambda & \sim \kappa^2 & \sim \lambda^2 & \sim \kappa^2 \\ \sim \lambda & \sim \lambda & \sim \lambda & \sim \kappa^2 & \sim \kappa^2 & \sim \lambda^2 \end{pmatrix}, \quad (2.9)$$

$$H_R = m_0^2 \begin{pmatrix} \sim \kappa^2 & \sim \kappa^2 & \sim \kappa^2 & 0 & \sim \kappa\lambda & \sim \kappa\lambda \\ \sim \kappa^2 & \sim \kappa^2 & \sim \kappa^2 & 0 & \sim \kappa\lambda & \sim \kappa\lambda \\ \sim \kappa^2 & \sim \kappa^2 & \sim \kappa^2 & 0 & \sim \kappa\lambda & \sim \kappa\lambda \\ 0 & 0 & 0 & * & * & * \\ \sim \kappa\lambda & \sim \kappa\lambda & \sim \kappa\lambda & * & \sim \lambda^2 & * \\ \sim \kappa\lambda & \sim \kappa\lambda & \sim \kappa\lambda & * & * & \sim \lambda^2 \end{pmatrix}. \quad (2.10)$$

Note that $(H_L)_{33} \ll (H_L)_{44}$, while $(H_R)_{33} \gg (H_R)_{44}$. This causes the exchange between the third and fourth rows in U_R in contrast to U_L . As a result, the mixing matrix U_R has matrix elements of the order

$$U_R = \begin{pmatrix} * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & * & * & * \\ * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & * & * & * \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & * & * & * \end{pmatrix}, \quad (2.11)$$

in contrast to

$$U_L = \begin{pmatrix} * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & * & * & * \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & * & * & * \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & * & * & * \end{pmatrix}. \quad (2.12)$$

The structures (2.11) and (2.12) mean that the dominant components of the fermions

in the up-quark sector are given by

$$\begin{aligned}
u_L &\simeq (2, 1) , & u_R &\simeq (1, 2) , \\
c_L &\simeq (2, 1) , & c_R &\simeq (1, 2) , \\
t_L &\simeq (2, 1) , & t_R &\simeq (1, 1) , \\
t'_L &\simeq (1, 1) , & t'_R &\simeq (1, 2) , \\
u_{5L} &\simeq (1, 1) , & u_{5R} &\simeq (1, 1) , \\
u_{6L} &\simeq (1, 1) , & u_{6R} &\simeq (1, 1) ,
\end{aligned} \tag{2.13}$$

of $SU(2)_L \times U(2)_R$, where we have denoted the fermion u_4 as t' . We should notice that t and t' have exceptional structures differently from other fermions f and F .

We consider that in the down-quark sector the seesaw expression (1.2) is well satisfied, so that the mixing matrices U_L and U_R are given by normal structure as (2.12). Then, the CKM matrix V_L is given by

$$V_L = \begin{pmatrix} * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ * & * & * & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 \\ \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \frac{1}{\lambda} & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 & \sim \left(\frac{1}{\lambda}\right)^2 \end{pmatrix}, \tag{2.14}$$

while the CKM matrix V_R for the right-handed weak currents is given by

$$V_R = \begin{pmatrix} * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 \\ * & * & * & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 \\ \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \frac{\kappa}{\lambda} & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 & \sim \left(\frac{\kappa}{\lambda}\right)^2 \end{pmatrix}, \tag{2.15}$$

where the factors $(1/\lambda)^2$ and $(\kappa/\lambda)^2$ come from the reason that the heavy fermions F_i are $(1, 1)$ of $SU(2)_L \times SU(2)_R$ in the present model.

III. CKM MATRICES V_L AND V_R

In order to see these relations (2.14) and (2.15) explicitly, we consider a

model with additional constraint

$$Z_R^T = Z_L, \quad Y^T = Y. \quad (3.1)$$

The constraint (3.1) are not so restrictive, and most seesaw mass matrix model will satisfy this constraint.

For the down-quark sector in which the seesaw expression (1.2) is valid, from (2.1) – (2.6), we obtain the relations

$$\begin{aligned} U_{dd}^R &\simeq (U_{dd}^L)^*, & U_{dD}^R &\simeq \kappa(U_{dD}^L)^*, \\ U_{Dd}^R &\simeq \kappa(U_{Dd}^L)^*, & U_{DD}^R &\simeq (U_{DD}^L)^*, \end{aligned} \quad (3.2)$$

where the 3×3 matrices U_{ab} ($a, b = f, F$) are defined by

$$U^f = \begin{pmatrix} U_{ff} & U_{fF} \\ U_{Ff} & U_{FF} \end{pmatrix}. \quad (3.3)$$

For the up-quark sector with $\det M_F = 0$, the seesaw expression (1.2) [therefore, (2.5) and (2.6)] is not valid any longer. However, when we define $\tilde{U}_R = P_{34}U_R$, where

$$P_{34} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (3.4)$$

we can see that the mixing matrix $\tilde{U}_R$ has a structure similar to U_L , because $\tilde{U}_R H_R \tilde{U}_R^\dagger = (DP_{34})^\dagger DP_{34} = \text{diag}(m_1^2, m_2^2, m_4^2, m_3^2, m_5^2, m_6^2)$ has the structure similar to $U_L H_L U_L^\dagger = D^2 = \text{diag}(m_1^2, m_2^2, m_3^2, m_4^2, m_5^2, m_6^2)$ apart from the exchange of the coefficients, $1 \leftrightarrow \kappa$ [see (2.9) and (2.10)]. Therefore, we obtain the relations

$$\begin{aligned} \tilde{U}_{uu}^R &\simeq (U_{uu}^L)^*, & \tilde{U}_{uU}^R &\simeq \kappa(U_{uU}^L)^*, \\ \tilde{U}_{Uu}^R &\simeq \kappa(U_{Uu}^L)^*, & \tilde{U}_{UU}^R &\simeq (U_{UU}^L)^*, \end{aligned} \quad (3.5)$$

similarly to (3.2).

Since the CKM mixing matrix V_L for the left-handed weak currents is given by

$$V_L = \begin{pmatrix} U_{uu}^L & U_{uU}^L \\ U_{Uu}^L & U_{UU}^L \end{pmatrix} \begin{pmatrix} \mathbf{1} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} U_{dd}^{L\dagger} & U_{Dd}^{L\dagger} \\ U_{dD}^{L\dagger} & U_{DD}^{L\dagger} \end{pmatrix} = \begin{pmatrix} U_{uu}^L U_{dd}^{L\dagger} & U_{uu}^L U_{Dd}^{L\dagger} \\ U_{Uu}^L U_{dd}^{L\dagger} & U_{Uu}^L U_{Dd}^{L\dagger} \end{pmatrix}, \quad (3.6)$$

the CKM mixing matrix V_R for the right-handed weak currents is given by

$$\begin{aligned} V_R &= P_{34} \begin{pmatrix} \tilde{U}_{uu}^R & \tilde{U}_{uU}^R \\ \tilde{U}_{Uu}^R & \tilde{U}_{UU}^R \end{pmatrix} \begin{pmatrix} \mathbf{1} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} U_{dd}^{R\dagger} & U_{Dd}^{R\dagger} \\ U_{dD}^{R\dagger} & U_{DD}^{R\dagger} \end{pmatrix} = P_{34} \begin{pmatrix} \tilde{U}_{uu}^R U_{dd}^{R\dagger} & \tilde{U}_{uu}^R U_{Dd}^{R\dagger} \\ \tilde{U}_{Uu}^R U_{dd}^{R\dagger} & \tilde{U}_{Uu}^R U_{Dd}^{R\dagger} \end{pmatrix} \\ &= P_{34} \begin{pmatrix} (V_{ud}^L)^* & \kappa(V_{uD}^L)^* \\ \kappa(V_{Ud}^L)^* & \kappa^2(V_{UD}^L)^* \end{pmatrix}. \end{aligned} \quad (3.7)$$

Therefore, we find

$$\begin{aligned} V_{ij}^R &= (V_{ij}^L)^* \quad i = u, c; \quad j = d, s, b, \\ V_{tj}^R &= (V_{t'j}^L)^* \quad j = d, s, b, \\ V_{t'j}^R &= (V_{tj}^L)^* \quad j = d, s, b. \end{aligned} \quad (3.8)$$

As seen from (3.8), the right-handed weak-interaction structure of t' is the same as the left-handed weak interaction structure of t .

In general, in the left-right symmetric model [7], the W_R -exchange diagrams can sizably contribute to the K^0 - $\bar{K}^0$ mixing [8]. However, in the present model, although the right-handed weak currents for u and c can contribute to the K^0 - $\bar{K}^0$ mixing as pointed out in Ref. [7], those for t and t' are negligibly small, because t_L (t'_R) is doublet of $SU(2)_{L(R)}$, while t_R (t'_L) is almost singlet of $SU(2)_{R(L)}$. Also the contributions for u_5 and u_6 are negligibly small because of the suppression factors $(1/\lambda)^2$ and $(\kappa/\lambda)^2$. For example, the K^0 - $\bar{K}^0$ mixing amplitude via (t, t', W_L, W_R) is suppressed by a factor $(1/\lambda)^2$ compared with that via (t, t, W_L, W_L) . The next leading term to the diagram (t, t, W_L, W_L) is a diagram (t', t', W_R, W_R) which is suppressed by a factor $(1/\kappa)^2$ compared with the diagram (t, t, W_L, W_L) . For $\kappa \geq 10$, the contributions are negligibly small.

Since the fourth up-quark t' has a comparatively light mass (of the order of m_{W_R}), we can expect the observation of t' -production via the reaction $d + u \rightarrow t' + d$ with the W_R exchange, for example, at LHC. Since we consider $m(W_R) \simeq \kappa m(W_L)$, we obtain

$$\sigma(p + p \rightarrow t' + X) \simeq \frac{1}{\kappa^4} \sigma(p + p \rightarrow t + X). \quad (3.9)$$

The decay width of t' , $\Gamma_{t'}$, is given by

$$\frac{\Gamma_{t'}}{\Gamma_t} \simeq \frac{m_{t'}^5/m_{W_R}^4}{m_t^5/m_{W_L}^4} \simeq \kappa , \quad (3.10)$$

from (3.8).

IV. NUMERICAL EXAMPLE FOR A SPECIFIC MODEL

Thus, as far as the mass matrix (1.1) satisfies the form (3.1), the CKM matrix V_R can be related to V_L irrelevantly to the explicit structures of Z and Y . However, in order to see the effects of the flavor-changing neutral currents (FCNC), we need the explicit forms of $U_{L(R)}^u$ and $U_{L(R)}^d$ separately, because the left- (right-) handed FCNC among the conventional fermions f_i ($i = 1, 2, 3$) are proportional [9] to the matrices (see Appendix)

$$C_{L(R)}^f = U_{fF}^{L(R)} (U_{fF}^{L(R)})^\dagger . \quad (4.1)$$

So, it is interesting to see the explicit structures of U_L and U_R for a realistic model which can give reasonable quark masses and the CKM mixing matrix parameters.

As an example, we choose the democratic seesaw mass matrix model [1], where M_F is given by the form [(unit matrix) + (a rank-one matrix)]:

$$M_F = \lambda m_0 Y_f = \lambda m_0 (\mathbf{1} + 3b_f X) , \quad (4.2)$$

where $\mathbf{1}$ and X are the 3×3 unit matrix and a rank-one matrix with the condition $X^2 = X$, respectively, and b_f is an f -dependent complex parameter. The name “democratic” [10] comes from the following assumption: the matrices Z_L and Z_R are given by a diagonal form in the heavy fermion basis on which the matrix X is democratic, i.e.,

$$X = \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} , \quad (4.3)$$

For simplicity, we assume that the matrices Z_L and Z_R have a common structure except for their phases:

$$Z_L = P(\delta_L) Z , \quad Z_R = P(\delta_R) Z , \quad (4.4)$$

$$P(\delta) = \text{diag}(e^{i\delta_1}, e^{i\delta_2}, e^{i\delta_3}) , \quad (4.5)$$

where the matrix Z is a real-parameter matrix

$$Z = \begin{pmatrix} z_1 & 0 & 0 \\ 0 & z_2 & 0 \\ 0 & 0 & z_3 \end{pmatrix} , \quad (4.6)$$

with $z_1^2 + z_2^2 + z_3^2 = 1$, and it is universal for all fermion sectors (up- and down-, quark and lepton sectors). In order to obtain input values for the parameters z_i , we assume that the parameter b_f takes the value $b_e = 0$ in the charged lepton sector, so that the parameters z_i are given by

$$\frac{z_1}{\sqrt{m_e}} = \frac{z_2}{\sqrt{m_\mu}} = \frac{z_3}{\sqrt{m_\tau}} = \frac{1}{\sqrt{m_e + m_\mu + m_\tau}} \quad (4.7)$$

from $M_e = m_0(\kappa/\lambda)P(\delta_L^e - \delta_R^e) \cdot Z \cdot \mathbf{1} \cdot Z$. The ansatz of the democratic M_F was motivated by the successful relation [11]

$$\frac{m_u}{m_c} \simeq \frac{3}{4} \frac{m_e}{m_\mu} , \quad (4.8)$$

(independently of κ/λ under $\lambda \gg \kappa$) for $b_u = -1/3$ and $b_e = 0$. In Ref. [1], the value of κ/λ has been fixed as $\kappa/\lambda = 0.02$ by the relations for $b_u \simeq -1/3$

$$m_c \simeq 2 \frac{m_\mu}{m_\tau} \frac{\kappa}{\lambda} m_0 , \quad m_t \simeq \frac{1}{\sqrt{3}} m_0 , \quad (4.9)$$

(note that the expression of m_t does not contain the suppression factor κ/λ). The value of $\beta_d \equiv \arg(-b_d)$ has been chosen as $\beta_d = \pi/10$ from the relations for $b_d \simeq -1$

$$m_s \simeq 2 \left| \sin \frac{\beta_d}{2} \right| \frac{m_\mu}{m_\tau} \frac{\kappa}{\lambda} m_0 , \quad m_b \simeq \frac{1}{2} \frac{\kappa}{\lambda} m_0 . \quad (4.10)$$

Then, we can find in Ref. [1] that these parameter values can successfully provide all of the quark mass ratios and CKM matrix parameters. It is worth while noting that the model can yield $m_t \gg m_b$ with keeping $m_u \sim m_d$ by adjusting only one complex parameter b_f and without choosing hierarchically different values between b_u and b_d .

In order to evaluate the CKM matrices V_L and V_R , it is convenient to define the matrix $\overline{M}$ which are given by

$$\overline{M} = P^\dagger(\delta_L) M P^\dagger(\delta_R) = m_0 \begin{pmatrix} 0 & Z \\ \kappa Z & \lambda Y \end{pmatrix}, \quad (4.11)$$

where the 6×6 phase matrix $P(\delta)$ is defined by

$$P(\delta) = \text{diag}(e^{i\delta_1}, e^{i\delta_2}, e^{i\delta_3}, 1, 1, 1), \quad (4.12)$$

(we have used the same notation with the 3×3 phase matrix (4.5)). The unitary matrices U_L and U_R are related to

$$U_L = \overline{U}_L P^\dagger(\delta_L), \quad U_R = \overline{U}_R P(\delta_R), \quad (4.13)$$

where

$$U_L M U_R^\dagger = \overline{U}_L \overline{M} \overline{U}_R^\dagger = D. \quad (4.14)$$

Then, the CKM matrices V_L and V_R are given by

$$\begin{aligned} V_L &= U_L^u P_0 U_L^{d\dagger} = \overline{U}_L^u P(\delta_L^d - \delta_L^u) P_0 \overline{U}_L^{d\dagger}, \\ V_R &= U_R^u P_0 U_R^{d\dagger} = \overline{U}_R^u P(\delta_R^u - \delta_R^d) P_0 \overline{U}_R^{d\dagger}, \end{aligned} \quad (4.15)$$

where

$$P_0 = \text{diag}(1, 1, 1, 0, 0, 0). \quad (4.16)$$

Since the constraint (3.1) means $\delta_{Ri}^f = \delta_{Li}^f \equiv \delta_i^f$, so that V_L and V_R are given by

$$V_L = \overline{U}_L^u P_0(\delta) \overline{U}_L^{d\dagger}, \quad V_R = \overline{U}_R^u P_0^\dagger(\delta) \overline{U}_R^{d\dagger}, \quad (4.17)$$

where

$$P_0(\delta) = \text{diag}(e^{i\delta_1}, e^{i\delta_2}, e^{i\delta_3}, 0, 0, 0), \quad (4.18)$$

with $\delta_i = -(\delta_i^u - \delta_i^d)$.

The observed CKM matrix parameters are roughly described by $(\delta_1, \delta_2, \delta_3) = (0, 0, \pi)$ [1] and more precisely by $(\delta_1, \delta_2, \delta_3) = (0, 0, \pi - \pi/30)$ [12]. However, in Refs. [1] and [12], only the 3×3 part of V_L has been investigated. Here, we show the numerical results of the 6×6 mixing matrices U_L and U_R for the case of $\kappa/\lambda = 0.02$, $\beta_d = \pi/10$, and $\delta_3 = \pi - \pi/30$ (we take $\kappa = 10$ temporarily according to Ref.[1],

but the results are almost insensitive to the value of κ):

$$U_L^u = \left(\begin{array}{ccc|ccc} +0.9994 & -0.0349 & -0.0084 & -0.0247\frac{1}{\lambda} & +6 \times 10^{-5}\frac{1}{\lambda} & +4 \times 10^{-6}\frac{1}{\lambda} \\ +0.0319 & +0.9709 & -0.2373 & -0.2051\frac{1}{\lambda} & -0.4345\frac{1}{\lambda} & +0.0259\frac{1}{\lambda} \\ +0.0165 & +0.2369 & +0.9714 & +0.8989\frac{1}{\lambda} & +0.8431\frac{1}{\lambda} & -0.0444\frac{1}{\lambda} \\ \hline +0.0093\frac{1}{\lambda} & +0.1114\frac{1}{\lambda} & -1.0364\frac{1}{\lambda} & +0.5774 & +0.5774 & +0.5772 \\ -0.0118\frac{1}{\lambda} & +0.1649\frac{1}{\lambda} & +0.0209\frac{1}{\lambda} & -0.7176 & +0.6961 & +0.0215 \\ -0.0064\frac{1}{\lambda} & -0.1011\frac{1}{\lambda} & +0.7927\frac{1}{\lambda} & -0.3894 & -0.4267 & +0.8163 \end{array} \right), \quad (4.19)$$

$$U_R^u = \left(\begin{array}{ccc|ccc} +0.9994 & -0.0349 & -0.0084 & -0.0247\frac{\kappa}{\lambda} & +6 \times 10^{-5}\frac{\kappa}{\lambda} & +4 \times 10^{-6}\frac{\kappa}{\lambda} \\ +0.0319 & +0.9709 & -0.2373 & -0.2051\frac{\kappa}{\lambda} & -0.4346\frac{\kappa}{\lambda} & +0.0259\frac{\kappa}{\lambda} \\ +0.0256\frac{\kappa}{\lambda} & +0.3459\frac{\kappa}{\lambda} & -0.0747\frac{\kappa}{\lambda} & +0.5773 & +0.5773 & +0.5774 \\ \hline +0.0165 & +0.2369 & +0.9713 & +0.3274\frac{\kappa}{\lambda} & +0.2716\frac{\kappa}{\lambda} & -0.6159\frac{\kappa}{\lambda} \\ -0.0118\frac{\kappa}{\lambda} & +0.1649\frac{\kappa}{\lambda} & +0.0209\frac{\kappa}{\lambda} & -0.7176 & +0.6961 & +0.0215 \\ -0.0064\frac{\kappa}{\lambda} & -0.1010\frac{\kappa}{\lambda} & +0.7929\frac{\kappa}{\lambda} & -0.3894 & -0.4267 & +0.8161 \end{array} \right). \quad (4.20)$$

$$|U_L^d| = \left(\begin{array}{ccc|ccc} 0.9772 & 0.2061 & 0.0506 & 0.0490\frac{1}{\lambda} & 0.0007\frac{1}{\lambda} & 4 \times 10^{-5}\frac{1}{\lambda} \\ 0.2118 & 0.9540 & 0.2124 & 0.2063\frac{1}{\lambda} & 0.0646\frac{1}{\lambda} & 0.0035\frac{1}{\lambda} \\ 0.0137 & 0.2179 & 0.9759 & 0.4335\frac{1}{\lambda} & 0.4809\frac{1}{\lambda} & 0.5251\frac{1}{\lambda} \\ \hline 0.0118\frac{1}{\lambda} & 0.1649\frac{1}{\lambda} & 0.0209\frac{1}{\lambda} & 0.7176 & 0.6961 & 0.0215 \\ 0.0064\frac{1}{\lambda} & 0.1010\frac{1}{\lambda} & 0.7927\frac{1}{\lambda} & 0.3895 & 0.4268 & 0.8162 \\ 0.0046\frac{1}{\lambda} & 0.0660\frac{1}{\lambda} & 0.2706\frac{1}{\lambda} & 0.5773 & 0.5773 & 0.5774 \end{array} \right), \quad (4.21)$$

where for U_L^d , for simplicity, we have shown only the magnitudes. Since the mixing matrix elements of U_R^d are given by the relation (3.5) with good approximation, here we have dropped the numerical result of U_R^d .

From (4.17), the 6×6 CKM matrix V_L is given by

$$|V_L| = \left(\begin{array}{ccc|ccc} 0.9756 & 0.2196 & 0.0028 & 0.0174\frac{1}{\lambda} & 0.0038\frac{1}{\lambda} & 0.0046\frac{1}{\lambda} \\ 0.2193 & 0.9749 & 0.0388 & 0.1615\frac{1}{\lambda} & 0.0910\frac{1}{\lambda} & 0.1283\frac{1}{\lambda} \\ 0.0105 & 0.0374 & 0.9992 & 0.0188\frac{1}{\lambda} & 0.7940\frac{1}{\lambda} & 0.2473\frac{1}{\lambda} \\ \hline 0.0780\frac{1}{\lambda} & 0.3253\frac{1}{\lambda} & 0.9873\frac{1}{\lambda} & 0.0399\left(\frac{1}{\lambda}\right)^2 & 0.8104\left(\frac{1}{\lambda}\right)^2 & 0.2879\left(\frac{1}{\lambda}\right)^2 \\ 0.0332\frac{1}{\lambda} & 0.1534\frac{1}{\lambda} & 0.0560\frac{1}{\lambda} & 0.0269\left(\frac{1}{\lambda}\right)^2 & 0.0331\left(\frac{1}{\lambda}\right)^2 & 0.0052\left(\frac{1}{\lambda}\right)^2 \\ 0.0626\frac{1}{\lambda} & 0.2638\frac{1}{\lambda} & 0.7517\frac{1}{\lambda} & 0.0331\left(\frac{1}{\lambda}\right)^2 & 0.6182\left(\frac{1}{\lambda}\right)^2 & 0.2212\left(\frac{1}{\lambda}\right)^2 \end{array} \right). \quad (4.22)$$

We have again dropped the results of V_R since the numerical results satisfies (3.7) very well.

The numerical results of C_L and C_R , to which the contributions of FCNC are proportional, are given as follows:

$$C_L^u = \begin{pmatrix} 2.43314 \times 10^{-9} & -2.013 \times 10^{-8} & -8.845 \times 10^{-8} \\ -2.013 \times 10^{-8} & 9.263 \times 10^{-7} & 2.208 \times 10^{-6} \\ -8.848 \times 10^{-8} & 2.208 \times 10^{-6} & 6.084 \times 10^{-6} \end{pmatrix}, \quad (4.23)$$

$$C_R^u = \begin{pmatrix} 2.43314 \times 10^{-7} & 2.013 \times 10^{-6} & 0.0002840 \\ 2.013 \times 10^{-6} & 9.264 \times 10^{-5} & 0.007087 \\ 0.0002840 & 0.007087 & 1.000 \end{pmatrix}, \quad (4.24)$$

$$|C_L^d| = \frac{1}{\kappa^2} |C_R^d| = \begin{pmatrix} 9.615 \times 10^{-9} & 4.026 \times 10^{-8} & 8.525 \times 10^{-8} \\ 4.026 \times 10^{-8} & 1.870 \times 10^{-7} & 3.514 \times 10^{-7} \\ 8.525 \times 10^{-8} & 3.514 \times 10^{-7} & 2.780 \times 10^{-6} \end{pmatrix}. \quad (4.25)$$

Thus, the matrix elements of C_L and C_R are suppressed by factors $(1/\lambda)^2$ and $(\kappa/\lambda)^2$, respectively, except for $(C_R^u)_{3i} = (C_R^u)_{i3}$ ($i = 1, 2, 3$). We see that the FCNC in the present model are harmless to the K^0 - $\bar{K}^0$ and D^0 - $\bar{D}^0$ mixings. However, the elements related to the top-quark have sizable values of C_R^u :

$$(C_R^u)_{tc} = 0.000709, \quad (C_R^u)_{tu} = 0.000284. \quad (4.26)$$

The observability of the single top-quark productions via FCNC with (4.26) will be discussed elsewhere [9].

V. SUMMARY

In conclusion, we have pointed out that in a seesaw quark mass matrix model which yields a singular enhancement of the top-quark mass, the 6×6 mixing matrix U_R^u for the right-handed up-quark sector has a peculiar structure, i.e., as if the third and fourth rows of U_R^u are exchanged in contrast to the left-handed mixing matrix U_L^u . This means that top quark t and the fourth up-quark t' have approximately components $t_L = (2, 1)$, $t_R = (1, 1)$, $t'_L = (1, 1)$ and $t'_R = (1, 2)$ of $SU(2)_L \times SU(2)_R$, although other fermions have $f_L = (2, 1)$, $f_R = (1, 2)$, $F_L = (1, 1)$ and $F_R = (1, 1)$.

For a model with the constraint (3.1) which is a likely case, the CKM mixing matrices V_L and V_R satisfy the relation (3.8). Observation of t' with mass $m_{t'} \simeq \kappa m_t$ (of the order of m_{W_R}) is expected at a future collider with a few TeV energy.

As an explicit example of U_L and U_R , we have investigated a model where M_F is given by a form (4.2). The numerical results, of course, satisfy the general relations (3.5) and (3.7). The matrix elements of $C = U_{fF}(U_{fF})^\dagger$ to which FCNC are proportional have been evaluated. The contributions of FCNC are harmless to the $K^0\text{-}\bar{K}^0$ and $D^0\text{-}\bar{D}^0$ mixings. On the other hand, the elements related to the top-quark have sizable values $(C_R^u)_{tc} = 0.000709$ and $(C_R^u)_{tu} = 0.000284$.

The present model which can successfully give quark mass ratios and CKM matrix parameters can also provide fruitful new physics. It seems that the model is worth investigating not only phenomenologically but also theoretically.

ACKNOWLEDGMENTS

The author would like to thank H. Fusaoka, T. Kurimoto and M. Tanimoto for helpful and enjoyable discussions. This work was supported by the Grant-in-Aid for Scientific Research, the Ministry of Education, Science and Culture, Japan (No.08640386).

APPENDIX: STRUCTURE OF FCNC

When the mass matrix M given in (1.1) is transformed as

$$\bar{\psi}_L M \psi_R + h.c. = \bar{\psi}'_L D \bar{\psi}'_R + h.c. \quad , \quad (A1)$$

where $\psi = (f, F)^T$, and $\psi' = U\psi$ is the mass-eigenstates, the vertex $\bar{\psi}_A \Gamma^{AB} \psi_B$ ($A, B = L, R$) is also transformed into $\bar{\psi}'_A \Gamma'^{AB} \psi'_B$, where

$$\Gamma'^{AB} = U_A \Gamma^{AB} U_B^\dagger \quad . \quad (A2)$$

For simplicity, hereafter, we drop the indices A, B . Correspondingly to (3.3), we denote the 6×6 matrix Γ in terms of 3×3 matrices Γ_{ab} ($a, b = f, F$) as

$$\Gamma = \begin{pmatrix} \Gamma_{ff} & \Gamma_{fF} \\ \Gamma_{Ff} & \Gamma_{FF} \end{pmatrix} \quad . \quad (A3)$$

Our interest is in the physical vertex Γ'_{ff} which is given by

$$\Gamma'_{ff} = \sum_a \sum_b U_{fa} \Gamma_{ab} U_{fb}^\dagger \quad , \quad (A4)$$

where $U_{ab}^\dagger \equiv (U_{ab})^\dagger = (U^\dagger)_{ba}$, because $(\Gamma'_{ff})_{ij}$ with $i \neq j$ mean transitions between f_i and f_j , i.e., appearance of the FCNC.

In our $SU(2)_L \times SU(2)_R \times U(1)_Y$ gauge model, the neutral currents $J_L^\mu = g_L^Z \bar{\psi} \Gamma_L^\mu \psi$, which couple with the left-handed weak boson Z_L^μ , are given by

$$\Gamma_L^\mu = \begin{pmatrix} c_L^f \mathbf{1} & 0 \\ 0 & c_L^F \mathbf{1} \end{pmatrix} \cdot \frac{1}{2} \gamma^\mu (1 - \gamma_5) + \begin{pmatrix} d_L^f \mathbf{1} & 0 \\ 0 & d_L^F \mathbf{1} \end{pmatrix} \cdot \frac{1}{2} \gamma^\mu (1 + \gamma_5) , \quad (A5)$$

where

$$\begin{aligned} c_L^f &= \pm \frac{1}{2} - \sin^2 \theta_L Q_f , \\ c_L^F &= -\sin^2 \theta_L Q_F , \end{aligned} \quad (A6)$$

$$\begin{aligned} d_L^f &= \pm \frac{1}{2} h_L - \sin^2 \theta_L Q_f , \\ d_L^F &= -\sin^2 \theta_L Q_F , \end{aligned} \quad (A7)$$

$$\sin^2 \theta_L = 1 - m_{W_L}^2 / m_{Z_L}^2 , \quad (A8)$$

$$h_L = -\frac{\sin^2 \theta_L}{1 - \varepsilon / \cos^2 \theta_L} \frac{\varepsilon}{\cos^2 \theta_L} , \quad (A9)$$

$$\varepsilon = m_{W_L}^2 / m_{W_R}^2 , \quad (A10)$$

the factor $\pm \frac{1}{2}$ takes $+\frac{1}{2}$ and $-\frac{1}{2}$ for up- and down-fermions, respectively, and Q_f (Q_F) is charge of the fermion f (F). Using the unitary condition for U_{ab} , $U_{ff} U_{ff}^\dagger + U_{fF} U_{fF}^\dagger = \mathbf{1}$, we can express the physical vertex Γ'_{Lff} as

$$\begin{aligned} \Gamma'_{Lff} &= \left(c_L^f U_{ff}^L U_{ff}^{L\dagger} + c_L^F U_{fF}^L U_{fF}^{L\dagger} \right) \cdot \frac{1}{2} \gamma^\mu (1 - \gamma_5) \\ &\quad + \left(d_L^f U_{ff}^R U_{ff}^{R\dagger} + d_L^F U_{fF}^R U_{fF}^{R\dagger} \right) \cdot \frac{1}{2} \gamma^\mu (1 + \gamma_5) \\ &= \left[c_L^f \mathbf{1} - (c_L^f - c_L^F) U_{fF}^L U_{fF}^{L\dagger} \right] \cdot \frac{1}{2} \gamma^\mu (1 - \gamma_5) \\ &\quad + \left[d_L^f \mathbf{1} - (d_L^f - d_L^F) U_{fF}^R U_{fF}^{R\dagger} \right] \cdot \frac{1}{2} \gamma^\mu (1 + \gamma_5) . \end{aligned} \quad (A11)$$

Similarly, for the neutral current $J_R^\mu = g_R^Z \bar{\psi} \Gamma_R^\mu \psi$, which couples with the right-handed weak boson Z_L , we obtain

$$\begin{aligned} \Gamma'_{Rff} &= \left[c_R^f \mathbf{1} - (c_R^f - c_R^F) U_{fF}^R U_{fF}^{R\dagger} \right] \cdot \frac{1}{2} \gamma^\mu (1 + \gamma_5) \\ &\quad + \left[d_R^f \mathbf{1} - (d_R^f - d_R^F) U_{fF}^L U_{fF}^{L\dagger} \right] \cdot \frac{1}{2} \gamma^\mu (1 - \gamma_5) , \end{aligned} \quad (A12)$$

where

$$\begin{aligned} c_R^f &= \pm \frac{1}{2} - \sin^2 \theta_R Q_f , \\ c_R^F &= -\sin^2 \theta_R Q_F , \end{aligned} \quad (A13)$$

$$\begin{aligned} d_R^f &= \pm \frac{1}{2} h_R - \sin^2 \theta_R Q_f , \\ d_R^F &= - \sin^2 \theta_R Q_F , \end{aligned} \quad (A14)$$

$$\sin^2 \theta_R = 1 - m_{W_R}^2 / m_{Z_R}^2 , \quad (A15)$$

$$h_R = - \frac{\sin^2 \theta_R}{1 - \varepsilon \cos^2 \theta_R} , \quad (A16)$$

$$\begin{aligned} g_R^Z &= -g_L^Z \frac{\sin \theta_L}{\sin \theta_R \cos \theta_R} \sqrt{\frac{1 - \varepsilon \cos^2 \theta_R}{1 - \varepsilon / \cos^2 \theta_L}} \\ &= \frac{e}{\cos \theta_L \sin \theta_R \cos \theta_R} \sqrt{\frac{1 - \varepsilon \cos^2 \theta_R}{1 - \varepsilon \cos^2 \theta_R / \cos^2 \theta_L}} . \end{aligned} \quad (A17)$$

Thus, the FCNC are induced by the second terms $U_{fF} U_{fF}^\dagger$ with magnitude $(c^f - c^F) [(d^f - d^F)]$. Therefore, we have denoted these matrices as $C_L^f \equiv U_{fF}^L U_{fF}^{L\dagger}$ and $C_R^f \equiv U_{fF}^R U_{fF}^{R\dagger}$ in (4.1).

References

- [1] Y. Koide and H. Fusaoka, Z. Phys. **C71**, 459 (1966).
- [2] T. Morozumi, T. Satou, M. N. Rebelo and M. Tanimoto, Preprint HUPD-9704 (1997), hep-ph/9703249.
- [3] Z. G. Berezhiani, Phys. Lett. **129B**, 99 (1983); Phys. Lett. **150B**, 177 (1985); D. Chang and R. N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987); A. Davidson and K. C. Wali, Phys. Rev. Lett. **59**, 393 (1987); S. Rajpoot, Mod. Phys. Lett. **A2**, 307 (1987); Phys. Lett. **191B**, 122 (1987); Phys. Rev. **D36**, 1479 (1987); K. B. Babu and R. N. Mohapatra, Phys. Rev. Lett. **62**, 1079 (1989); Phys. Rev. **D41**, 1286 (1990); S. Ranfone, Phys. Rev. **D42**, 3819 (1990); A. Davidson, S. Ranfone and K. C. Wali, Phys. Rev. **D41**, 208 (1990); I. Sogami and T. Shinohara, Prog. Theor. Phys. **66**, 1031 (1991); Phys. Rev. **D47**, 2905 (1993); Z. G. Berezhiani and R. Rattazzi, Phys. Lett. **B279**, 124 (1992); P. Cho, Phys. Rev. **D48**, 5331 (1994); A. Davidson, L. Michel, M. L. Sage and K. C. Wali, Phys. Rev. **D49**, 1378 (1994); W. A. Ponce, A. Zepeda and R. G. Lozano, Phys. Rev. **D49**, 4954 (1994).

- [4] M. Gell-Mann, P. Rammond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, 1979); T. Yanagida, in *Proc. Workshop of the Unified Theory and Baryon Number in the Universe*, edited by A. Sawada and A. Sugamoto (KEK, 1979); R. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [5] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **73**, 225 (1994).
- [6] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1996); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [7] J. C. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974); R. N. Mohapatra and J. C. Pati, Phys. Rev. **D11**, 366 and 2588 (1975); G. Senjanovic and R. N. Mohapatra, Phys. Rev. **D12**, 1502 (1975).
- [8] T. Kurimoto, A. Tomita and S. Wakaizumi, Phys. Lett. **B381**, 470 (1996).
- [9] Y. Koide, Preprint US-96-09 (1996), hep-ph/9701261 (unpublished).
- [10] H. Terazawa, University of Tokyo, Report No. INS-Rep.-298 (1977) (unpublished); H. Harari, H. Haut and J. Weyers, Phys. Lett. **78B**, 459 (1978); T. Goldman, in *Gauge Theories, Massive Neutrinos and Proton Decays*, edited by A. Perlmutter (Plenum Press, New York, 1981), p.111; T. Goldman and G. J. Stephenson, Jr., Phys. Rev. **D24**, 236 (1981); Y. Koide, Phys. Rev. Lett. **47**, 1241 (1981); Phys. Rev. **D28**, 252 (1983); **39**, 1391 (1989); C. Jarlskog, in *Proceedings of the International Symposium on Production and Decays of Heavy Hadrons*, Heidelberg, Germany, 1986 edited by K. R. Schubert and R. Waldi (DESY, Hamburg), 1986, p.331; P. Kaus, S. Meshkov, Mod. Phys. Lett. **A3**, 1251 (1988); Phys. Rev. **D42**, 1863 (1990); L. Lavoura, Phys. Lett. **B228**, 245 (1989); M. Tanimoto, Phys. Rev. **D41**, 1586 (1990); H. Fritzsch and J. Plankl, Phys. Lett. **B237**, 451 (1990); Y. Nambu, in *Proceedings of the International Workshop on Electroweak Symmetry Breaking*, Hiroshima, Japan, (World Scientific, Singapore, 1992), p.1.
- [11] Y. Koide, Mod. Phys. Lett. **A8**, 2071 (1993).
- [12] Y. Koide and H. Fusaoka, Prog. Theor. Phys. **97**, 459 (1966).